

SPECIAL VALUES OF CANONICAL GREEN'S FUNCTIONS

ROBIN DE JONG

ABSTRACT. We give a precise formula for the value of the canonical Green's function at a pair of Weierstrass points on a hyperelliptic Riemann surface. Further we express the 'energy' of the Weierstrass points in terms of a spectral invariant recently introduced by N. Kawazumi and S. Zhang. It follows that the energy is strictly larger than $\log 2$. Our results generalize known formulas for elliptic curves.

1. INTRODUCTION

Let M be a compact and connected Riemann surface of genus $h \geq 1$. On M one has a canonical Kähler form μ given as follows. The space H of holomorphic differentials on M is a complex vector space of dimension h . It carries a natural hermitian inner product given by $\langle \alpha, \beta \rangle = \frac{\sqrt{-1}}{2} \int_M \alpha \wedge \bar{\beta}$ for elements α, β in H . Let $(\omega_1, \dots, \omega_h)$ be an orthonormal basis of H . By the Riemann-Roch theorem, the ω_i do not simultaneously vanish at any point of M , and we put

$$\mu = \frac{\sqrt{-1}}{2h} \sum_{i=1}^h \omega_i \wedge \bar{\omega}_i.$$

The associated metric has been studied in many contexts, ranging from arithmetic geometry [3] [9] [10] [19] [21] to perturbative string theory [1] [2] [5] [6] [13]. The metric is known to have everywhere non-positive curvature, and the curvature vanishes at $x \in M$ if and only if x is a Weierstrass point on a hyperelliptic Riemann surface [20], Main Theorem.

In this paper we study the canonical Green's function g_μ associated to μ (see Section 2 for definitions), and in particular its special values $g_\mu(w_i, w_j)$ at pairs (w_i, w_j) of distinct Weierstrass points on a hyperelliptic Riemann surface. The starting point of our discussion is the case $h = 1$, with M given as a 2-sheeted cover of the Riemann sphere $\mathbb{CP}^1$ branched at four points, say $\alpha_1, \alpha_2, \alpha_3$ and ∞ . Let w_1, w_2, w_3, o be the four critical points of M lying over $\alpha_1, \alpha_2, \alpha_3, \infty$. It is then known [7] [15] [22] that the formula

$$(1.1) \quad g_\mu(w_i, w_j) = \frac{1}{3} \log 2 + \frac{1}{12} \log \left(\frac{|\alpha_i - \alpha_j|^2}{|\alpha_i - \alpha_k| \cdot |\alpha_j - \alpha_k|} \right)$$

holds. By the translation invariance of g_μ we obtain the formula

$$(1.2) \quad \sum_{i=1}^3 g_\mu(w_i, o) = \log 2.$$

2010 *Mathematics Subject Classification*. Primary 14H55; secondary 14H45, 14H15.

Key words and phrases. Canonical Green's function, hyperelliptic curves, Kawazumi-Zhang invariant, Weierstrass points.

We will return to both these formulas in the text below.

Our purpose is to generalize formulas (1.1) and (1.2) to the case of a hyperelliptic Riemann surface of genus $h \geq 2$. Let M be such a surface, given as the 2-sheeted cover of $\mathbb{CP}^1$ branched at the $2h+2$ points $\alpha_1, \dots, \alpha_{2h+2}$ (these may or may not include ∞). We put, for each pair (i, j) of distinct indices,

$$(1.3) \quad \delta_{ij} = \frac{(\alpha_i - \alpha_j)^{2h(2h+1)} \prod_{r \neq s} (\alpha_r - \alpha_s)}{\prod_{r \neq i} (\alpha_i - \alpha_r)^{2h+1} \prod_{r \neq j} (\alpha_j - \alpha_r)^{2h+1}}.$$

In this formula, we disregard any difference of two roots when one of the roots is ∞ . It is readily verified that δ_{ij} is invariant under the action of $\text{Aut}(\mathbb{CP}^1) = \text{PGL}_2(\mathbb{C})$ on the α_i . In particular δ_{ij} defines an analytic modular invariant of the pair (w_i, w_j) , where w_i, w_j are the critical points of M lying over α_i, α_j . More intrinsically δ_{ij} is the discriminant of the branch set that one obtains by sending α_i, α_j to $0, \infty$ using an element of $\text{Aut}(\mathbb{CP}^1)$ and by normalizing the other $2h$ roots such that their product is unity [12].

We note that there is a tight connection with the more familiar cross ratios on the α_i . Put $\eta_{ijk} = \delta_{ik} \delta_{jk}^{-1}$, and let

$$\mu_{ijk} = (\alpha_i - \alpha_k)(\alpha_j - \alpha_r)(\alpha_j - \alpha_k)^{-1}(\alpha_i - \alpha_r)^{-1}$$

be the cross ratio of the 4-tuple $(\alpha_i, \alpha_j, \alpha_k, \alpha_r)$. Then the relation

$$\mu_{ijk}^{2h(2h+1)} = \eta_{ijk} \eta_{ijr}^{-1} = \delta_{ik} \delta_{jk}^{-1} \delta_{jr} \delta_{ir}^{-1}$$

holds. It is straightforward to verify that for each index i , the product $\prod_{j \neq i} \delta_{ij}$ is equal to unity.

The *energy* of the Weierstrass points on M is defined to be the following invariant of M :

$$\psi(M) = \frac{1}{2h+2} \sum_{i \neq j} g_\mu(w_i, w_j),$$

where the summation runs over all pairs i, j of distinct indices. Our first result is the following.

Theorem A. *Let w_i, w_j be two distinct Weierstrass points of M . Then the formula*

$$g_\mu(w_i, w_j) = \frac{1}{4h(2h+1)} \log |\delta_{ij}| + \frac{1}{2h+1} \psi(M)$$

holds.

In our second result we give an expression for $\psi(M)$. Consider for the moment an arbitrary compact and connected Riemann surface M of genus $h \geq 1$. Let Δ_μ be the Laplacian on $L^2(M, \mu)$ given by putting $\partial \bar{\partial} f = \pi \sqrt{-1} \Delta_\mu(f) \mu$ for C^∞ functions in $L^2(M, \mu)$. Let $(\phi_\ell)_{\ell=0}^\infty$ be an orthonormal basis of real eigenfunctions of Δ_μ , with corresponding eigenvalues $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots$. We then let $\varphi(M)$ denote the invariant

$$\varphi(M) = \sum_{\ell > 0} \frac{2}{\lambda_\ell} \sum_{m, n=1}^h \left| \int_M \phi_\ell \omega_m \wedge \bar{\omega}_n \right|^2$$

of (M, μ) . This fundamental invariant was introduced and studied recently by N. Kawazumi [18] and S. Zhang [24]. Note that $\varphi(M) \geq 0$, and that $\varphi(M) = 0$ if and only if M has genus one. In [17] the asymptotic behavior of the φ -invariant is determined for surfaces degenerating into a stable curve with a single node. The

result shows in particular that φ can be viewed as a Weil function, with respect to the boundary, on the stable (Deligne-Mumford) compactification of the moduli space of surfaces of a fixed genus $h \geq 2$.

Theorem B. *Let M be a hyperelliptic Riemann surface of genus $h \geq 2$, and let $\varphi(M)$ be the Kawazumi-Zhang invariant of M . Then for the energy $\psi(M)$ of the set of Weierstrass points of M , the formula*

$$\psi(M) = \frac{1}{2h} \varphi(M) + \log 2$$

holds.

In particular we find that $\psi(M) > \log 2$. With $h = 1$, the formulas in Theorems A and B specialize to equations (1.1) and (1.2), respectively.

In order to prove Theorem B we proceed as follows. First we prove that, when viewed as functions on the moduli space of hyperelliptic Riemann surfaces, both ψ and $\frac{1}{2h}\varphi$ have the same image under $\partial\bar{\partial}$. Then we study the asymptotic behavior of the difference of ψ and $\frac{1}{2h}\varphi$ on a family of hyperelliptic Riemann surfaces degenerating into a stable curve of compact type. By using induction on h we will then deduce that the difference is a constant on the moduli space, equal to $\log 2$. For the degeneration argument we will rely heavily on results obtained by R. Wentworth [23].

2. CANONICAL GREEN'S FUNCTIONS

In this section we collect some basic results on canonical Green's functions. The main sources are [3] [10]. Proposition 2.2, about a 'parallel' property of certain $(1, 1)$ -forms related to the universal Riemann surface over the moduli space of Riemann surfaces, seems not to be well known, and is possibly of independent interest.

As before, let M be a compact and connected Riemann surface of genus $h \geq 1$ and μ its canonical Kähler form. Let Δ be the diagonal on the complex manifold $M \times M$. The canonical Green's function $g_\mu = g_\mu(x, y)$ is the C^∞ -function on $M \times M \setminus \Delta$ uniquely characterized by the following conditions:

- (1) $\partial\bar{\partial}g_\mu(x, y) = \pi\sqrt{-1}(\mu(y) - \delta_x)$ for all $x \in M$;
- (2) $\int_M g_\mu(x, y)\mu(y) = 0$ for all $x \in M$;
- (3) $g_\mu(x, y) = g_\mu(y, x)$ for all $x \neq y \in M$;
- (4) $g_\mu(x, y) - \log|z(x) - z(y)|$ is bounded for all $x \neq y$ in a coordinate chart $(U, z: U \xrightarrow{\sim} D)$ of M , where D is the open unit disk.

The canonical Green's function inverts the Laplacian Δ_μ in the sense that

$$f(x) = - \int_M g_\mu(x, y) \Delta_\mu(f)(y) \mu(y) + \int_M f \mu$$

holds for all $x \in M$ and all f in $C^\infty(M)$. As a consequence we have a formal development

$$g_\mu(x, y) = \sum_{\ell > 0} \frac{\phi_\ell(x)\phi_\ell(y)}{\lambda_\ell}$$

for g_μ , where ϕ_ℓ and λ_ℓ are the eigenfunctions and eigenvalues of Δ_μ that we have introduced above.

In [10] Section 7 using standard harmonic analysis explicit formulas are derived for the ϕ_ℓ and λ_ℓ in the case $h = 1$. As a corollary of these formulas [10] Section 7

contains a derivation of equation (1.2) from the resulting formal expression for g_μ . We would like to present here an alternative argument.

Proposition 2.1. *Assume (M, o) is a complex torus, and let N be a positive integer. Let $x_1, \dots, x_{N^2-1}$ be the non-trivial N -torsion points of M . Then the formula*

$$\sum_{i=1}^{N^2-1} g_\mu(x_i, o) = \log N$$

holds.

Proof. From properties (1) and (2) it follows that for every x, y on M with $Nx \neq y$ the equality

$$g_\mu(Nx, y) = \sum_{w: Nw=y} g_\mu(x, w)$$

holds. Now choose x in a standard euclidean coordinate chart $z: U \xrightarrow{\sim} D$ around o , where D is the open unit disk. Then we have

$$\begin{aligned} g_\mu(Nx, o) &= \sum_{w: Nw=o} g_\mu(x, w) \\ &= \sum_{i=1}^{N^2-1} g_\mu(x_i, x) + g_\mu(x, o) \\ &= \sum_{i=1}^{N^2-1} g_\mu(x_i, x) + \log |z(x)| + a + o(1) \end{aligned}$$

as $x \rightarrow o$, where a is some constant, by properties (3) and (4). On the other hand we have

$$\begin{aligned} g_\mu(Nx, o) &= \log |Nz(x)| + a + o(1) \\ &= \log N + \log |z(x)| + a + o(1) \end{aligned}$$

as $x \rightarrow o$, by property (4). The desired equality follows. $\square$

One obtains (1.2) by taking $N = 2$ in the above proposition.

Let $\mathcal{M}$ denote the moduli space of compact and connected Riemann surfaces of genus $h \geq 2$, and denote by $\pi: \mathcal{C} \rightarrow \mathcal{M}$ the universal surface over $\mathcal{M}$. Both are viewed as orbifolds. The canonical Green's functions on the fibers of π determine a generalized function g on the fiber product $\mathcal{C} \times_{\mathcal{M}} \mathcal{C}$ with logarithmic singularities along the diagonal Δ . Let ν be the $(1, 1)$ -form on $\mathcal{C} \times_{\mathcal{M}} \mathcal{C}$ determined by the condition $\partial\bar{\partial}g = \pi\sqrt{-1}(\nu - \delta_\Delta)$. Let e^A denote the restriction of ν to Δ , which we view as another copy of $\mathcal{C}$. By [3] the form e^A restricts to $2 - 2h$ times the canonical $(1, 1)$ -form μ in each fiber of $\pi: \mathcal{C} \rightarrow \mathcal{M}$.

The $(1, 1)$ -forms e^A and ν are parallel over $\mathcal{M}$ in the following sense.

Proposition 2.2. *Let s, t be arbitrary holomorphic sections of π over an open subset U of $\mathcal{M}$. Then the equalities $s^*e^A = t^*e^A = (s, t)^*\nu$ hold on U .*

Proof. Without loss of generality we may replace $\mathcal{M}$ by a finite cover so that we may assume that $\mathcal{M}$ is equipped with a universal theta characteristic α , i.e. a consistent choice of a divisor class α of degree $h - 1$ on each surface M such that 2α is the canonical class. Let $\mathcal{J} \rightarrow \mathcal{M}$ be the universal jacobian over $\mathcal{M}$ and let $\Phi: \mathcal{C} \times_{\mathcal{M}} \mathcal{C} \rightarrow \mathcal{J}$ be the map over $\mathcal{M}$ given by sending a triple (M, x, y) to the pair

consisting of the jacobian $J(M)$ of M and the class of the divisor $h \cdot x - y - \alpha$ in $J(M)$. Let $\bar{\pi}: \mathcal{C} \times_{\mathcal{M}} \mathcal{C} \rightarrow \mathcal{C}$ be the projection on the first factor and let $\bar{s}, \bar{t}: \pi^{-1}U \rightarrow \mathcal{C} \times_{\mathcal{M}} \mathcal{C}$ be the local sections of $\bar{\pi}$ obtained by pulling back the given local sections s, t of π . It follows from [14] Lemma 3.2 that there exists a $(1, 1)$ -form κ on $\mathcal{C}$ and a $(1, 1)$ -form w on $\mathcal{J}$ which is translation-invariant in the fibers of $\mathcal{J} \rightarrow \mathcal{M}$, such that $\Phi^*w = h\nu + \bar{\pi}^*\kappa$ holds. We deduce from this that

$$h \bar{s}^* \nu = (\Phi \bar{s})^* w - \kappa \quad \text{and} \quad h \bar{t}^* \nu = (\Phi \bar{t})^* w - \kappa.$$

Let $T_{x,y}$ denote translation by $[x - y]$ over $\mathcal{J}|_U$. Then $\Phi \bar{t} = T_{s,t} \Phi \bar{s}$ and since w is fiberwise translation-invariant we obtain $\bar{s}^* \nu = \bar{t}^* \nu$ from the above equalities. We derive $(s, t)^* \nu = s^* \bar{t}^* \nu = s^* \bar{s}^* \nu = (s, s)^* \nu = s^* e^A$. By symmetry property (3) we have $(s, t)^* \nu = (t, s)^* \nu = t^* e^A$ as well. $\square$

3. PROOF OF THEOREM A

In this section we give a proof of Theorem A. Let $x: M \rightarrow \mathbb{CP}^1$ be the 2-sheeted cover with branch points $\alpha_1, \dots, \alpha_{2h+2}$. Let $w_1, \dots, w_{2h+2}$ be the corresponding Weierstrass points on M . For the moment we fix two distinct indices i, j . Consider then the meromorphic function $f_{ij} = (x - \alpha_i)(x - \alpha_j)^{-1}$ on M . We note that $\text{div}(f_{ij}) = 2(w_i - w_j)$. From properties (1) and (2) we therefore obtain an equality

$$g_\mu(w_i, z) - g_\mu(w_j, z) = \frac{1}{2} \log |f_{ij}(z)| - \frac{1}{2} \int_M \log |f_{ij}| \cdot \mu$$

of generalized functions on M . In particular we find for any pair of indices k, r

$$\begin{aligned} g_\mu(w_i, w_k) - g_\mu(w_i, w_r) - g_\mu(w_j, w_k) + g_\mu(w_j, w_r) \\ &= \frac{1}{2} \log |f_{ij}(w_k) f_{ij}(w_r)^{-1}| \\ &= \frac{1}{2} \log |\mu_{ijk r}| \\ &= \frac{1}{4h(2h+1)} \log |\eta_{ijk} \eta_{ijr}^{-1}| \end{aligned}$$

where we recall that $\eta_{ijk} = \delta_{ik} \delta_{jk}^{-1}$ with δ_{ij} the expression from (1.3). This equality implies that there exists a constant c_{ij} such that

$$g_\mu(w_i, w_k) - g_\mu(w_j, w_k) = \frac{1}{4h(2h+1)} \log |\eta_{ijk}| + c_{ij}$$

for each index k . It follows from [16], Theorem 1.4 that

$$\sum_{k \neq i} g_\mu(w_i, w_k) = \sum_{k \neq j} g_\mu(w_j, w_k).$$

From the fact that $\prod_{k \neq i, j} \delta_{ij}$ equals unity we obtain that $\sum_{k \neq i, j} \log |\eta_{ijk}| = 0$. It follows that

$$2h c_{ij} = \sum_{k \neq i, j} (g_\mu(w_i, w_k) - g_\mu(w_j, w_k)) = 0,$$

hence $c_{ij} = 0$ and

$$g_\mu(w_i, w_k) - g_\mu(w_j, w_k) = \frac{1}{4h(2h+1)} \log |\eta_{ijk}| = \frac{1}{4h(2h+1)} \log |\delta_{ik} \delta_{jk}^{-1}|.$$

Now fix the index k and vary the indices i, j . We find that there exists a constant c_k such that

$$g_\mu(w_i, w_k) = \frac{1}{4h(2h+1)} \log |\delta_{ik}| + c_k$$

for all indices $i \neq k$. We obtain $c_k = \frac{1}{2h+1} \psi(M)$ by summing over $i \neq k$. The formula in Theorem A follows.

Note that a similar argument in the case $h = 1$ allows one to deduce (1.1) from (1.2).

4. PROOF OF THEOREM B

As was announced in the Introduction, we proceed in several steps. Let $\mathcal{H}$ be the moduli space of hyperelliptic Riemann surfaces of a fixed genus $h \geq 2$. Let $\pi: \mathcal{C} \rightarrow \mathcal{H}$ be the universal surface over $\mathcal{H}$. Both are viewed as orbifolds. A first result is the following.

Proposition 4.1. *The equality of $(1, 1)$ -forms*

$$\partial \bar{\partial} \psi = \frac{1}{2h} \partial \bar{\partial} \varphi$$

holds on $\mathcal{H}$.

Proof. Let $\mathcal{U}$ be an open cover of $\mathcal{H}$ such that for each U in $\mathcal{U}$ there exist $2h+2$ holomorphic sections $w_1, \dots, w_{2h+2}: U \rightarrow \pi^{-1}U$ of $\pi^{-1}U \rightarrow U$ such that each w_i is a Weierstrass point in each fiber of $\pi^{-1}U \rightarrow U$. Let U be an element of $\mathcal{U}$. It suffices to prove the required equality on U . Theorem 3.1 of [18] gives an expression for $\partial \bar{\partial} \varphi$ over the moduli space of Riemann surfaces of genus h (note that [18] considers the invariant a with $a = \frac{1}{2\pi} \varphi$). Since the ‘harmonic volume’ of a Weierstrass point on a hyperelliptic Riemann surface is trivial, this expression immediately implies that on U one has the equality

$$\partial \bar{\partial} \varphi = 2h(2h+1)\pi\sqrt{-1}w_i^*e^A,$$

for any choice of the index i . On the other hand by Proposition 2.2 we have

$$\partial \bar{\partial} \psi = \pi\sqrt{-1} \sum_{j \neq i} (w_i, w_j)^* \nu = (2h+1)\pi\sqrt{-1}w_i^*e^A$$

for each index i . The proposition follows. $\square$

We next determine the limit behavior of the difference $\psi - \frac{1}{2h} \varphi$ in a holomorphic family M_t of hyperelliptic Riemann surfaces of genus h degenerating into a stable curve M_0 which is the union of two hyperelliptic Riemann surfaces M_1, M_2 of genera $h_1, h_2 \geq 1$, respectively, with two points $p_1 \in M_1$ and $p_2 \in M_2$ identified, as in Chapter III of the book ‘Theta functions on Riemann surfaces’ by J. Fay [11]. Here t runs through a small punctured open disk around 0 in the complex plane. Note that $h = h_1 + h_2$. Let g_1, g_2 be the canonical Green’s functions on M_1, M_2 , respectively, and let z_1, z_2 be the local coordinates around p_1, p_2 on M_1, M_2 that come with the degeneration model. We put, following [23], Section 6:

$$\log k_1 = \lim_{x \rightarrow p_1} [g_1(x, p_1) - \log |z_1(x)|], \quad \log k_2 = \lim_{x \rightarrow p_2} [g_2(x, p_2) - \log |z_2(x)|],$$

and then make the reparametrization $\tau = k_1 k_2 t$.

Theorem 4.2. *For the surfaces M_t in Fay's degeneration model, the limit formula*

$$\lim_{t \rightarrow 0} \left[\frac{1}{2h} \varphi(M_t) + \frac{h_1 h_2}{h^2} \log |\tau| \right] = \frac{1}{2h} \varphi(M_1) + \frac{1}{2h} \varphi(M_2)$$

holds.

Theorem 4.3. *For the energy $\psi(M_t)$ of the Weierstrass points of M_t , the limit formula*

$$\lim_{t \rightarrow 0} \left[\psi(M_t) + \frac{h_1 h_2}{h^2} \log |\tau| \right] = \frac{h_1}{h} \psi(M_1) + \frac{h_2}{h} \psi(M_2)$$

holds.

We deduce Theorem B from Theorems 4.2 and 4.3 by induction on h . Note that the assertion in Theorem B is true for $h = 1$ by (1.2) and the fact that φ vanishes for $h = 1$. Now assume that the assertion in Theorem B is true for all genera smaller than h . Consider a holomorphic family M_t of hyperelliptic Riemann surfaces of genus h as above. We find

$$\begin{aligned} \lim_{t \rightarrow 0} \left[\psi(M_t) - \frac{1}{2h} \varphi(M_t) \right] &= \frac{h_1}{h} \psi(M_1) - \frac{1}{2h} \varphi(M_1) + \frac{h_2}{h} \psi(M_2) - \frac{1}{2h} \varphi(M_2) \\ &= \frac{h_1}{h} \left(\psi(M_1) - \frac{1}{2h_1} \varphi(M_1) \right) + \frac{h_2}{h} \left(\psi(M_2) - \frac{1}{2h_2} \varphi(M_2) \right) \\ &= \frac{h_1}{h} \log 2 + \frac{h_2}{h} \log 2 = \log 2. \end{aligned}$$

Let $\overline{\mathcal{H}}$ be the closure of $\mathcal{H}$ in the Deligne-Mumford compactification $\overline{\mathcal{M}}$ of the moduli space $\mathcal{M}$ of compact Riemann surfaces of genus h . Let $\tilde{\mathcal{H}} \subset \overline{\mathcal{H}}$ be the partial compactification of $\mathcal{H}$ given by stable curves of compact type (i.e. whose jacobians are principally polarized abelian varieties). Write $\chi = \psi - \frac{1}{2h} \varphi$. The above calculation implies that for hyperelliptic Riemann surfaces in $\mathcal{H}$ tending to a point in the boundary of $\mathcal{H}$ in $\tilde{\mathcal{H}}$, the function χ tends to $\log 2$. In particular the function χ extends as a continuous function over $\tilde{\mathcal{H}}$. By Proposition 4.1 we know that χ is a pluriharmonic function on $\mathcal{H}$. By the Riemann extension theorem we find that χ extends as a pluriharmonic function on $\tilde{\mathcal{H}}$, which is a constant function with value $\log 2$ on $\tilde{\mathcal{H}} \setminus \mathcal{H}$.

Let $\mathcal{A}^*$ be the Satake compactification of the moduli space $\mathcal{A}$ of principally polarized abelian varieties of genus h . This is a normal projective variety [4]. There exists a holomorphic map $t: \overline{\mathcal{H}} \rightarrow \mathcal{A}^*$ given by assigning to a stable curve M the jacobian of its normalization. The map t is injective on $\mathcal{H}$, and the function χ descends as a pluriharmonic function on $t(\tilde{\mathcal{H}})$, which is a constant function with value $\log 2$ on $t(\tilde{\mathcal{H}} \setminus \mathcal{H})$.

The boundary of $\tilde{\mathcal{H}}$ in $\overline{\mathcal{H}}$ is a finite union of divisors Ξ_i , see [8] for example. From the moduli theoretic description of the points in the divisors Ξ_i it follows that the map t restricted to the boundary has positive dimensional fibers. Hence the boundary of $t(\tilde{\mathcal{H}})$ in the closure of $t(\mathcal{H})$ in $\mathcal{A}^*$ has codimension at least two. Let M be a point in $t(\mathcal{H})$. Using that $\mathcal{A}^*$ is projective one sees from intersecting with suitable hyperplanes in a projective embedding that the variety $t(\tilde{\mathcal{H}})$ contains a complete curve X passing through M . As $t(\mathcal{H})$ is affine, the curve X necessarily contains a point of $t(\tilde{\mathcal{H}} \setminus \mathcal{H})$. Let $Y \rightarrow X$ be the normalization of X . The function

χ pulls back to a pluriharmonic function on Y , which is then necessarily constant. It follows that the value of χ at M is equal to $\log 2$, and Theorem B is proven.

For the proof of Theorem 4.2 we refer to [17]. It remains to prove Theorem 4.3. We need the next result, which follows from [23], Theorem 6.10.

Theorem 4.4. *Let g_t be the canonical Green's function on M_t . Denote by p both the point p_1 on M_1 and the point p_2 on M_2 . Then for local distinct holomorphic sections x, y of the family M_t specializing onto M_1 we have*

$$\lim_{t \rightarrow 0} \left[g_t(x, y) - \left(\frac{h_2}{h} \right)^2 \log |\tau| \right] = g_1(x, y) - \frac{h_2}{h} (g_1(x, p) + g_1(y, p)) .$$

For local distinct holomorphic sections x, y with x specializing onto M_1 and y specializing onto M_2 we have

$$\lim_{t \rightarrow 0} \left[g_t(x, y) + \frac{h_1 h_2}{h^2} \log |\tau| \right] = \frac{h_1}{h} g_1(x, p) + \frac{h_2}{h} g_2(y, p) .$$

Now let $w_1, \dots, w_{2h+2}$ be the Weierstrass points on M_t . As $t \rightarrow 0$ a portion of $2h_1 + 1$ of them degenerate onto M_1 , and $2h_2 + 1$ of them degenerate onto M_2 . The point p is a Weierstrass point of both M_1, M_2 . Let I be the set of indices such that $i \in I$ if and only if w_i degenerates onto M_1 .

Lemma 4.5. *The equality*

$$\sum_{\substack{i, j \in I \\ i \neq j}} g_1(w_i, w_j) = 2h_1 \psi(M_1)$$

holds.

Proof. We calculate

$$(2h_1 + 2) \psi(M_1) = \sum_{\substack{i, j \in I \cup \{p\} \\ i \neq j}} g_1(w_i, w_j) = \sum_{\substack{i, j \in I \\ i \neq j}} g_1(w_i, w_j) + 2 \psi(M_1) .$$

The required equality follows. $\square$

Using Theorem 4.4 we find

$$\begin{aligned} (2h + 2) \psi(M_t) &= \sum_{i \neq j} g_t(w_i, w_j) \\ &= \sum_{\substack{i, j \in I \\ i \neq j}} g_t(w_i, w_j) + \sum_{\substack{i, j \in I^c \\ i \neq j}} g_t(w_i, w_j) + 2 \sum_{\substack{i \in I \\ j \in I^c}} g_t(w_i, w_j) \\ &= \sum_{\substack{i, j \in I \\ i \neq j}} \left(\left(\frac{h_2}{h} \right)^2 \log |\tau| + g_1(w_i, w_j) - \frac{h_2}{h} (g_1(w_i, p) + g_1(w_j, p)) \right) \\ &\quad + \sum_{\substack{i, j \in I^c \\ i \neq j}} \left(\left(\frac{h_1}{h} \right)^2 \log |\tau| + g_2(w_i, w_j) - \frac{h_1}{h} (g_2(w_i, p) + g_2(w_j, p)) \right) \\ &\quad + 2 \sum_{\substack{i \in I \\ j \in I^c}} \left(-\frac{h_1 h_2}{h^2} \log |\tau| + \frac{h_1}{h} g_1(w_i, p) + \frac{h_2}{h} g_2(w_j, p) \right) + o(1) \end{aligned}$$

as $t \rightarrow 0$. With the help of Lemma 4.5 this can be rewritten as

$$\begin{aligned}
(2h+2)\psi(M_t) &= \left((2h_1+1)2h_1 \left(\frac{h_2}{h}\right)^2 + (2h_2+1)2h_2 \left(\frac{h_1}{h}\right)^2 \right) \log |\tau| \\
&\quad - 2(2h_1+1)(2h_2+1) \frac{h_1 h_2}{h^2} \log |\tau| \\
&\quad + \left(2h_1 - \frac{4h_1 h_2}{h} + \frac{2h_1(2h_2+1)}{h} \right) \psi(M_1) \\
&\quad + \left(2h_2 - \frac{4h_1 h_2}{h} + \frac{2h_2(2h_1+1)}{h} \right) \psi(M_2) + o(1) \\
&= (2h+2) \left(-\frac{h_1 h_2}{h^2} \log |\tau| + \frac{h_1}{h} \psi(M_1) + \frac{h_2}{h} \psi(M_2) \right) + o(1)
\end{aligned}$$

as $t \rightarrow 0$. Theorem 4.3 follows.

REFERENCES

- [1] L. Alvarez-Gaumé, J.-B. Bost, G. Moore, P. Nelson, C. Vafa, *Bosonization on higher genus Riemann surfaces*. Comm. Math. Phys. 112 (1987), 503–552.
- [2] L. Alvarez-Gaumé, G. Moore, C. Vafa, *Theta functions, modular invariance, and strings*. Comm. Math. Phys. 106 (1986), 1–40.
- [3] S.Y. Arakelov, *An intersection theory for divisors on an arithmetic surface*. Izv. Akad. USSR 86 (1974), 1164–1180.
- [4] W.L. Baily, A. Borel, *On the compactification of arithmetically defined quotients of bounded symmetric domains*. Bull. Amer. Math. Soc. 70 (1964) 588–593.
- [5] A.A. Beilinson, Y.I. Manin, *The Mumford form and the Polyakov measure in string theory*. Comm. Math. Phys. 107 (1986), 359–376.
- [6] A.A. Belavin, V.G. Knizhnik, *Complex geometry and the theory of quantum strings*. Soviet Phys. JETP 64 (1986), 214–228.
- [7] M. Bershadsky, A. Radul, *Fermionic fields on $\mathbb{Z}_N$ -curves*. Comm. Math. Phys. 116 (1988), no. 4, 689–700.
- [8] E. Cornalba, J. Harris, *Divisor classes associated to families of stable varieties, with applications to the moduli space of curves*. Ann. Sci. Ec. Norm. Sup. 21 (1988), 455–475.
- [9] J.-M. Couveignes, B. Edixhoven (eds.), *Computational aspects of modular forms and Galois representations*. Annals of Mathematics Studies 176, Princeton University Press 2011.
- [10] G. Faltings, *Calculus on arithmetic surfaces*. Ann. of Math. 119 (1984), 387–424.
- [11] J. Fay, *Theta functions on Riemann surfaces*. Lecture Notes in Mathematics 352, Springer Berlin Heidelberg New York, 1973.
- [12] J. Guàrdia, *Jacobian Nullwerte, periods and symmetric equations for hyperelliptic curves*. Ann. de l'Inst. Fourier 57 no. 4 (2007), 1253–1283.
- [13] E. d'Hoker, H.D. Phong, *The geometry of string perturbation theory*. Rev. Modern Phys. 60 (1988), 917–1065.
- [14] R. de Jong, *Arakelov invariants of Riemann surfaces*. Doc. Math. 10 (2005), 311–329.
- [15] R. de Jong, *On the Arakelov theory of elliptic curves*. Enseign. Math. 51 (2005), 179–201.
- [16] R. de Jong, *Faltings' delta-invariant of a hyperelliptic Riemann surface*. In: G. van der Geer, B. Moonen, R. Schoof (eds.), *Number Fields and Function Fields – Two Parallel Worlds*. Progress in Mathematics vol. 239, Birkhäuser Verlag 2005.
- [17] R. de Jong, *Asymptotic behavior of the Kawazumi-Zhang invariant for degenerating Riemann surfaces*. Preprint, [arXiv:1207.2353](https://arxiv.org/abs/1207.2353).
- [18] N. Kawazumi, *Johnson's homomorphisms and the Arakelov-Green function*. Preprint, [arXiv:0801.4218](https://arxiv.org/abs/0801.4218).
- [19] S. Lang, *Introduction to Arakelov theory*. Springer Verlag, Berlin Heidelberg New York, 1988.
- [20] D. Lewittes, *Differentials and matrices on Riemann surfaces*. Trans. Amer. Math. Soc. 139 (1969), 311–318.

- [21] C. Soulé, *Géométrie d'Arakelov des surfaces arithmétiques*. Séminaire Bourbaki 1988/89. Astérisque 177-178 (1989), Exp. 713, 327–343.
- [22] L. Szpiro, *Sur les propriétés numériques du dualisant relatif d'une surface arithmétique*. In: The Grothendieck Festschrift, Vol. III, 229–246, Progr. Math. 88, Birkhäuser, Boston, 1990.
- [23] R. Wentworth, *The asymptotics of the Arakelov-Green's function and Faltings' delta invariant*. Comm. Math. Phys. 137 (1991), 427–459.
- [24] S. Zhang, *Gross-Schoen cycles and dualizing sheaves*. Invent. Math. 179 (2010), 1–73.

Address of the author:

Mathematical Institute
University of Leiden
PO Box 9512
2300 RA Leiden
The Netherlands
Email: `rdejong@math.leidenuniv.nl`